LIST OF FIGURES

<u>Figure</u>		<u>Page</u>
I-1.	Annual total reconstructed light extinction coefficient (Mm ⁻¹) calculated from the aerosol concentrations at each operational site in the IMPROVE network for the period March 1992 through February 1995	I-20
I-2.	Seasonal visibility across the continental United States calculated using b _{ext} data from each operational transmissometer site in the IMPROVE network for the period March 1988 through February 1995	I-21
I-3	Annual average percentage of total light extinction (including Rayleigh) contributed by sulfate, nitrate, organics, elemental carbon, and coarse particles for the central Rocky Mountain, northern Rocky Mountain, and northern Great Plains regions (March 1988 through February 1995).	
I-4.	Annual average measured fine aerosol mass budgets (in percent) for the central Rocky Mountains, northern Rocky Mountains, and northern Great Plains regions (March 1988 through February 1995).	
II-1.	National parks and major cities of the Rocky Mountain and northern Great Plains regions	
II-2.	Air quality monitoring and meteorological sites for Colorado	II-7
II-3.	Air quality monitoring and meteorological sites for Wyoming	
II-4.	Air quality monitoring and meteorological sites for Montana	
II-5.	Air quality monitoring and meteorological sites for North Dakota	
II-6.	Air quality monitoring and meteorological sites for South Dakota	
II-7.	1996 SO ₂ point sources for the Rocky Mountain and northern Great Plains regions	
II-8.	1996 NO _x point sources for the Rocky Mountain and northern Great Plains regions	
III-1.	Soils map of the Loch Vale watershed (from Walthall 1985	
III-2.	Vegetation classification map of ROMO, from park database	
III-3.	Map of ROMO and air quality monitoring sites	
III-4.	Diurnal ozone concentrations at ROMO and other locations along the Front Range on July 2, 1993	
III-5.	Nitrate concentration measured in alpine lakes east of the Continental Divide	
	throughout the length of Colorado and in southeastern Wyoming	III-26
III-6.	Watersheds of ROMO studied by the U.S. Fish and Wildlife Service	III-27
III-7.	Predicted acidity (expressed as pH) of snowmelt at Buffalo Pass, adjacent to the Mt. Zirkel Wilderness Area, compared with expected salamander egg mortality at	
	various pH values	III-37
III-8.	Daily discharge (A) and nitrate concentration (B) in Icy Brook and Andrews Creek within the Loch Vale watershed in April-September 1992	III-39
III-9.	Seasonal average reconstructed extinction (Mm ⁻¹) for the period March 1988 through February 1995	III-46
III-10.	Reconstructed extinction budgets for Rocky Mountain National Park, Colorado, March 1988 - February 1995	III-48
III-11.	Annual arithmetic mean and cumulative frequency statistics, Rocky Mountain National Park, Colorado, transmissometer data (filtered)	
III-12.	Photographs illustrating visibility conditions at Rocky Mountain National Park	III-54
IV-1.	Soil classifications of GRTE	\/-2
IV-1.	Land cover of GRTE	
IV-2.	The elevational distribution of major forest types near GRTE	
IV-4.	Location of lakes sampled by Gulley and Parker in GRTE	
V-1.	Regional-geological map of YELL (adapted from Cox 1973 by Gibson et al. 1983)	
V-2.	Major vegetation types, roads, and large lakes in YELL	
V-3.	Air quality monitoring locations at YELL	

<u>Figure</u>		<u>Page</u>
V-4. V-5.	Alkalinity map for lakes in YELL, prepared from existing data by Gibson et al. (1983). Seasonal average reconstructed extinction (Mm ⁻¹) YELL, March 1988 through	V-17
	February 1995	V-20
V-6.	Reconstructed extinction budgets for YELL, March 1988 - February 1995	V-21
V-7.	Annual arithmetic mean and cumulative frequency statistics for YELL,	
	transmissometer data (filtered)	V-26
V-8.	Photographs illustrating visibility conditions at Yellowstone National Park	V-28
VI-1.	Annual average daily fluoride emissions from Columbia Falls Aluminum Co. plant	
	from 1957 to 1992 (Colombia Falls Aluminum Co., Columbia Falls, MT)	VI-6
VI-2.	Location of the air quality monitoring site at GLAC.	VI-9
VI-3.	Mean alkalinity concentrations regressed against mean conductivity values for the	1/1/4/4
\ /I \ 4	frontcountry and backcountry lakes of GLAC, 1984-1990. R ² =0.989	VI-11
VI-4.	Time series plot of pH in Snyder, Upper Dutch and Cobalt Lakes, GLAC,	V/I 40
\/I E	1984-1990Transects of vegetation sampling for fluoride done in Flathead NF and GLAC in	V I-12
VI-5.	1970 by USDA Forest Service and National Air Pollution Control Administration	\/ 17
VI-6.	Isolines of fluoride (ppb) measured in foliage samples taken along transects from	V I- I /
v 1-0.	Columbia Falls Aluminum Company in 1970 (provided by GLAC)	\/I_1 Q
VI-7.	Permanent vegetation sampling locations in the southwestern portion of the park	
VI-7.	Seasonal average reconstructed extinction (Mm ⁻¹) GLAC, March 1988 through	VI-13
V1 O.	February 1995	VI-21
VI-9.	Reconstructed extinction budgets for GLAC, Montana, March 1988 - February 1995	
VI-10.	Annual arithmetic mean and cumulative frequency statistics, GLAC, Montana,	0
	transmissometer data (filtered).	VI-27
VI-11	Photographs illustrating visibility conditions at Glacier National Park	
VII-1.	Oil and gas well locations in the vicinity of the North Unit	
VII-2.	Oil and gas well locations in the vicinity of the South Unit	VII-9
VII-3.	Water quality monitoring locations in THRO	VII-17
VII-4.	Water quality monitoring site identification numbers in the North Unit (Panel A) and	
	the South Unit (Panel B)	VI-18
VII-5.	Measured Ca ²⁺ , Mg ²⁺ , and pH values of water quality monitoring sites for which	
	data are available	
VIII-1.	Distribution of forest and grassland vegetation in WICA.	VIII-3
VIII-2.		
	WICA	VIII-7
IX-1.	Seasonal average reconstructed extinction (Mm ⁻¹) for BADL, March 1988 through	D4 4 =
13.4.5	February 1995.	
IX-2.	Reconstructed extinction budgets for BADL, March 1988 - February 1995	IX-11
IX-3.	Annual arithmetic mean and cumulative frequency statistics, BADL, South Dakota,	11/ 45
17/ 4	transmissometer data (filtered)	
IX-4.	Photographs illustrating visibility conditions at Badlands National Park	IX-1/